



DISINFECTANT SPRAY

- CLEANER
- DEODORIZER
- DISINFECTANT
- *VIRUCIDE
- TUBERCULOCIDAL
- MILDEWSTAT
- FUNGICIDE
- Citrus Scent
- Ready-To Use Disinfectant Spray
- Eliminates Odors
- Cleans and Deodorizes in One Easy Step
- Kills 99.9% of Germs and Bacteria

ACTIVE INGREDIENTS


n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS	99.790%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panels for additional precautionary statements.

EPA REG. NO. 1839-83-83894
EPA EST. NO. 83894-LA-001

 Distributed by Q-Based Solutions, Inc.[™]
A Division of Q-Based Healthcare
Erath, LA 70533 1.337.937.8800
Info@Qbased.com www.Qbased.com

NET CONTENTS - 32oz (946 mL)
U.S. Patent No. 5,444,094

PuraCleanRx Disinfectant Spray is specifically designed as a general cleaner and disinfectant for use in homes, nursing homes, schools, locker rooms, schools, transportation facilities, manufacturing facilities, lodging facilities, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities and where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces.

DISINFECTION, DEODORIZING AND CLEANING - Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for **HIV-1 1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

One-step cleaning and disinfection of toilet bowls and urinals: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of PuraCleanRx Disinfectant Spray evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet. For HIV-1 **1 minute** and for HBV and HCV **5 minutes**) and flush.

Disinfection of animal quarters and kennels: For disinfection of pre-cleaned animal quarters and kennels, apply PuraCleanRx Disinfectant Spray. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **3 minutes**; for Bovine Viral Diarrhea Virus (BVDV) **5 minutes**; for Feline Calicivirus and Rabies virus **30 seconds**; for Canine Parvovirus **10 minutes**. Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Using approved AOAC test methods under Good Laboratory Practices, [GLPs], in the presence of 5% blood serum and a **3 minute** contact time, unless otherwise noted, PuraCleanRx Disinfectant Spray kills the following organisms on hard non-porous inanimate surfaces:

BACTERICIDAL ACTIVITY: Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, Escherichia coli O157:H7, Streptococcus pyogenes, Listeria monocytogenes, Yersinia enterocolitica, Enterococcus faecium, Corynebacterium ammoniagenes, Salmoneella typhi

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), Vancomycin intermediate resistant Staphylococcus aureus (VISA), (CDC Isolate 99287), Methicillin resistant Staphylococcus epidermidis (MRSE)

***VIRUCIDAL ACTIVITY -** PuraCleanRx Disinfectant Spray kills on hard, non-porous inanimate surfaces when allowed to remain wet for a **10 minute** contact time against: Hepatitis A Virus (HAV), Canine Parvovirus, Poliovirus Type1

PuraCleanRx Disinfectant Spray, when used on environmental, inanimate hard surfaces exhibits virucidal activity against **Hepatitis B Virus (HBV)**, **Hepatitis C Virus (HCV)** and **Bovine Viral Diarrhea Virus (BVDV)** when the treated surface is allowed to remain wet for **5 minutes**; for **Human Coronavirus** and **SARS associated Coronavirus, 2 minutes**; for **HIV-1 (associated with AIDS), 1 minute**.

KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

CONTACT TIME: Allow surface to remain wet for HBV and HCV 5 minutes, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

FUNGICIDAL ACTIVITY - PuraCleanRx Disinfectant Spray is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet.

MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION KEEP OUT OF REACH OF CHILDREN.

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

if in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

if swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

if inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

Store in a dry place no lower in temperature than 50 F or higher than 120 F. Do not reuse empty container. Wrap container and put in trash.

Container Disposal for Refill Spray Bottle: Do not reuse empty spray bottle except with refill bottle, otherwise discard in trash. Do not reuse empty refill bottle. Wrap refill bottle and discard in trash.